

Global Journal of Health Science (GJHS)

PROTECTIVE ROLE OF HERBAL PLANTS IN HEPATOTOXICITY

S.Babyvanitha, Vinothkumar and B.Jaykar



Protective Role of Herbal Plants in Hepatotoxicity



S.Babyvanitha¹, Vinothkumar², B.Jaykar³

Corresponding Author's Email: info@iprjb.org

Article History

Received 10th January 2023

Received in Revised Form 25th January 2023

Accepted 10th February 2023



Abstract

Purpose: The liver is largest metabolic organ involved in detoxification and synthesis of essential biomolecules in the human body. Liver damage occurs due to chronic alcohol abuse, viral infections (Hepatitis A, B, C) medicines, poor hygiene, smoking, dietary substances, chemicals (Carbon tetra chloride, thioacetamide, acetaminophen, D-Galactosamine) or herb induced hepatotoxicity. The liver filters all the blood from the digestive tract, before passing it to the rest of the body to avoid the entry of toxins in the other body system and prevent severe consequences. Hepatic insufficiency may increase plasma total cholesterol and triacylglycerols with high risk of atherosclerosis and cardiovascular disease. Free radical formation in the alcohol use result in development of hepatitis leading to cirrhosis. Plants have been used to protect liver injury by various chemicals.

Methodology: In this study we reviewed some popular herbal plants having hepatoprotective potential.

Findings: Polyphenols, flavonoids, terpenoids are natural antioxidants and the consumption of foods that contain these compounds in large quantities seems to play an important role in prophylaxis against many diseases.

Unique Contribution to Theory, Practice and Policy: Herbal drugs are safe and have potential to cure hepatotoxicity. In India many plants have been identified as hepatoprotective drugs and used to treat liver disorders.

Keywords: *Herbal Plants, Hepatotoxicity, Liver*

©2023 by the Authors. This Article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>)

INTRODUCTION

Liver is major metabolic organ in human body and also involved in excretion of many endogenous and exogenous compounds¹. An injury to liver or impairment of its function may develop many consequences on individual health². Liver damage is associated with increase in tissue lipid peroxidation and depletion of reduced glutathione levels. Serum level of many biochemical markers such as transaminases, alkaline phosphatase, bilirubin, triglycerides and cholesterol are elevated in liver disease. Liver involved in the metabolism of nutrients like carbohydrates, proteins, lipids and excretion of waste metabolites and detoxification of exogenous and endogenous compounds like xenobiotics, drugs, viral infections and chronic alcoholism. A liver disease is a worldwide problem. Conventional drugs used in the treatment of liver diseases are inadequate and have some serious side effects. Numerous medicinal plants and their formulations are used for liver disorders in ethnomedical practice as well as traditional system of medicine in India. Many naturally occurring products have been reported to contain large amount of antioxidant other than vitamin C, E and carotenoid.

Liver-Anatomy

Liver is the largest gland in the body weighing about 1-2.3 kg. Located in the upper part of the abdominal cavity occupying right hypochondriac region, part of the epigastric region and extending into the left hypochondriac region. Liver is enclosed in thin inelastic capsule and incompletely covered by peritoneum. Peritoneal folds form supporting ligaments that attach liver to interior surface of the diaphragm. The liver has four lobes; right lobe, left lobe, caudate and quadrate lobes. Through portal fissure various structures enter and leave the gland; the portal vein enters carrying blood from the stomach, spleen, pancreas and small and large intestines. The hepatic artery enters carrying arterial blood. Sympathetic and parasympathetic nerves enter into the liver through this fissure. The right and left hepatic duct leaves carrying bile from the liver to the gallbladder. Lymph vessels leave the liver draining lymph to abdominal and thoracic nodes. The lobes of liver are made up of small functional units called lobules. It is formed by cuboidal cells, the hepatocytes arranged in pairs of columns radiating from a central vein. Between two pairs of column of cells are sinusoids containing mixture of blood from the portal vein and hepatic artery. It allows mixing arterial and portal venous blood and coming into close contact with the liver cells. The cells lining the sinusoids are kupffer cells that ingest and destroy worn out blood cells and foreign particles present in the blood flowing through the liver. Blood drains from the sinusoids into centrilobular veins then merge with veins of other lobules forming larger veins leave the liver and empty into the inferior vena cava.

Functions of the Liver

The liver regulates plasma glucose level. After meal blood glucose level increases, while insulin from pancreas convert the glucose into glycogen and stored in liver. When the blood glucose level falls glucagon from pancreas convert the glycogen into glucose and maintain the normal blood glucose level. Liver metabolise fat to a form in which it can be used by the tissues to provide energy. The liver involved in deamination of amino acids- removes the nitrogenous portion from aminoacids that are not required for protein synthesis; urea is formed from the nitrogenous part and is excreted in urine. It remove nitrogenous portion of amino acid and attaches it to other carbohydrate molecules forming new non-essential amino acids. Plasma proteins are formed from amino acids includes albumin, globulins and blood clotting factors

by the liver. Kupffer cells in the sinusoids breakdown the red blood cells. Liver produces immune factors that fight against the microbes. Phagocytes in the liver produce acute-phase proteins in response to microbes. These proteins are associated with inflammation process, tissue repair and immune cell activities. Liver also involved in fat metabolism i.e. oxidation of fatty acids, formation of lipoproteins, cholesterol and phospholipids. Fatty acids formed by the digestion of lipids used to synthesize cholesterol. It acts as a storage site for vitamin A, K, B12, and D, iron. The liver also stores fats, copper⁴.

Removing and excreting body wastes, hormones and other foreign substances. These substances enter the blood through production by metabolism within the body or from outside in the form of drugs or foreign compounds. Breakdown of red blood cells liberate haemoglobin which lead to formation of bilirubin and is excreted in to bile by hepatocytes. If bilirubin cannot be removed from the blood due to gallstones, liver disease or excessive breakdown of red blood cells produce jaundice⁵. Bile formation is helpful in digestion. Bile is secreted by liver and stored in gallbladder and it enters into the duodenum of the small intestine.

Pathophysiology of Hepatotoxicity

Metabolism and close relationship with the gastrointestinal tract, the liver is susceptible to injury from drugs and other substances. Seventy five percent of the blood coming to the liver arrives directly from gastrointestinal organ and then spleen via portal vein that brings drugs and xenobiotics. Many chemicals damage mitochondria, an organelle that produces energy. Mitochondrial impairment releases more amounts of oxidants that injure hepatic cells. Activation of CYP2E1 leads to oxidative stress which plays a crucial role in liver damage. Injury to hepatocyte and bile duct cells leads to accumulation of bile acid inside the liver. This promotes further liver damage.

Patterns of Injury

Chemicals produce a clinical and pathological hepatic injury. Biochemical markers such as alanine transferase, alkaline phosphatase and bilirubin used to indicate liver damage. Liver injury is defined as a rise in either ALT level more than three times of the upper limit of normal (ULN). Alkaline phosphatase (ALP) level more than twice ULN when associated with increased ALT or ALP⁶.The liver synthesizes concentrates and secretes bile acid and excretes other toxicants such as bilirubin. Drug induced injury to hepatocytes and bile duct cells can lead to chloestasis. It leads to intrahepatic accumulation of toxic bile acids and excretion products which promote further hepatic injury⁷.

Marker enzymes such as ALP, AST, and ALT used to assess the liver damage. Membrane damage or necrosis releases the enzyme in to circulation that can be measured in the serum. High level of AST indicates liver damage caused by viral hepatitis, cardiac infarction and muscle injury. AST catalyses conversion of alanine to pyruvate and glutamate. ALT is better parameter for detecting liver injury. Elevated level of serum enzymes are indicative of cellular leakage and loss of functional integrity of cell membrane in liver⁸.

APAP Induced Hepatotoxicity

Adverse hepatic events caused by drugs may be either predictable or unpredictable. Paracetamol produce predictable liver injury occur within few days as a result of direct liver toxicity of the parent drug or its metabolites. Unpredictable events occur as overt or symptomatic disease within 1-8 weeks like phenytoin or longer one year period of latency such

as isoniazid. Most of adverse drug- induced hepatic events are unpredictable and are either immune mediated hypersensitivity reactions or idiosyncratic. The pathogenesis of drug induced hepatotoxicity occurs by toxic drug or metabolite that either elicits an immune response or directly affects the biochemistry of the cell. In both case the resultant cell death is the event that leads to the clinical manifestation hepatitis. Metabolism of chemicals takes place in liver

Acetaminophen is widely used as analgesic and antipyretic for mild to moderate pain and fever. Harmless at low doses, but has direct hepatotoxic potential when taken as an overdose and cause acute liver injury and cell death. Even in therapeutic doses acetaminophen cause serum aminotransferase elevations. Acetaminophen is converted to nontoxic glucuronate or sulphate conjugates and secreted in the urine. A minor amount of acetaminophen is metabolized by CYP450 to N-acetyl-p-benzoquinoneimine that can be toxic. Normally this intermediate is fastly conjugated to reduce glutathione, detoxified and secreted. If level of glutathione are low or the pathway is overwhelmed by high doses of acetaminophen, the reactive intermediate accumulates and bind to intracellular macromolecules that can lead to cell injury through apoptotic pathway. Factors that increase the metabolism of acetaminophen through CYP450 or that decrease the availability of glutathione can predispose to acetaminophen toxicity. Factors that affect downstream toxicity of acetaminophen metabolic intermediates may also affect toxicity. These factors are important in designing therapies for acetaminophen hepatotoxicity.

Table 1: Ethnomedicinal Plants Used in Liver Diseases

Plant name	Family	Chemical constituents	Traditional uses	References
<i>Artemisia scoparia</i>	Compositae	Hyperin, eupafolin, pedalitin, 5,7,2', 4'-tetrahydroxy-6,5'-dimethoxy flavones, camphor, and 1,8-, and beta-caryophyllene	Jaundice and liver disorders	9,10,11
<i>Apium graveolens</i> Linn	Apiaceae	Limonene, p-dimethyl styrene, n-pertyl benene, caryophyllene, a-selinene, n-butyl phthalide, and sedanenolide	Liver and spleen disorders, jaundice, rheumatism, gout and inflammatory diseases	12-14, 15
<i>Allium hirtifolium</i>	Alliaceae	Saponins, sapogenins, thiosulfinate, shallomin, quercetin, Kaempferol(12), Alliin, alliinase, allicin, s-allyl-cysteine, diallyl disuophide, diallyltrisulphide, Methylallyltrisulphide(13)	Fatty acids-9-hexadecenoic acid, 14-eicosadienoic acid, n-hexadecanoic acid, linolenic acid, linoleic acid	14
<i>Apium graveolens</i>	Umbelliferae	Isoimperatorin, isoquercetin, linoleic acid, coumarin, furanocoumarins, apigenin, apiin, phenolic compounds, choline and unidentified alkaloids. It contains betacarotene, folic acid, vitamin C, sodium, magnesium, silica, potassium, Chlorophyll and fiber. Essential oil contains deltalimonene and various sesquiterpene.	Antifungal, antihypertensive, antihyperlipidemic, diuretic, anticancer, antioxidative, hepatoprotective	16,17,18,19
<i>Cynara scolymus</i>	Astraceae	Cynarin, luteolin, chlorogenic acid, caffeic acid, flavonoids, polyphenol	Used for digestive disorders, hyperlipidemia, liver and bile diseases, hepatoprotective, antioxidant, antimicrobial	20, 21
<i>Berberis Vulgaris</i>	Berberidaceae	Berberine, oxyacanthine, berbamine, palmatine, columbamine, malic acid, jatrorrhizine, berberrubine, stigmasterol, terpenoids, lupeol, oleanolic acid, stigmasterol glucoside, polyphenols	Antioxidant, hepatoprotective agents.	22,23,24
<i>Calandula officinalis</i>	Asteraceae	A-thujene, T-murolol, sesquiterpene, flavonol glycosides, triterpene alcohols, triterpenoid saponins, flavonoids, carotenoids, xanthophylls, phenolic acids, mucilage, bitters, phytosterols, tocopherols, calendulin, resin, and volatile oils.	Antibacterial, anti-inflammatory, antiviral, antioxidant, antipyretics, jaundice, Hepatoprotective activity	25,26,27
<i>Nigella Sativa</i>	Ranunculaceae	Thymoquinone, p-cymene, α -pinene,	Asthma, cough, bronchitis, headache, rheumatism, fever, influenza, eczema, anti-inflammatory	28, 29, 30, 31
<i>Taraxacum officinale</i>	Asteraceae	Sesquiterpene, lactones, triterpenes, carbohydrates, fattyacids, carotenoids, flavonoids, minerals, taraxalisin, coumarins and cichoriin, aesculin, Germacrane, caffeic acid, chlorogenic acid, luteolin and luteolin-7-glucoside. α -pinene, β -pinene	Antioxidant, Anti-inflammatory, Hepatoprotective.	32, 33
<i>Allium Sativum</i>	Amaryllidaceae	Sapogenins, saponins, sulphuric compounds and flavonoids, S-alkylcysteine, S-allylmercaptocysteine, (alpha) – fructosyl arginine, allicin	Antidiabetic, anti-inflammatory, antihypertension, antimicrobial, antiatherosclerotic, and hepatoprotective	34,35
<i>Marrubium vulgare</i>	Lamiaceae	Diterpene, sterols, Phenylpropanoids, flavonoids, (E)-caryophyllene, germacrene D, bicyclogermacrene	Antihypertensive, analgesic, anti-inflammatory, hypoglycaemic effect, antidyplipidemic effect and antioxidant	36,37
<i>Ammi majus</i>	Apiaceae	Quercetin, Kaempferol, marmesinin	Psoriasis, vitiligo, Diabetic nephropathy, myocardial injury	107
<i>Citrullus lanatus</i>	Curcubitaceae	Cucurbitain, triterpenes, anthraquinones, sterols, alkaloids, flavonoids, saponins, tannins, flavones aglycone, and phenols.	Cytoprotective, antioxidant, antifibrosis	38,39
<i>Agrimonia eupatoria</i>	Rosaceae	Phenolic compounds, coumarin, flavonoid, tannins, terpenoids, procatechuic acid, coumaric acid, chlorogenic acid, quercitin, gallic acid	Hepatoprotective activity	40
<i>Prunus armeniaca</i>	Rosaceae	Salicylic acid, tannins, potassium salts, pcoumaric acid, procatechuic acid, ferulic acid and diferulic acids. 4'.5,7-trihydroxy flavones-7-O(β -D-mannopyranosyl, β -D-galactopyranoside. B-D-allopyranoside and 3, 4', 5, 7 –tetrahydroxy-3.5'-di-methoxy flavones 3-O-, carotenoids, flavonoids, phenols.	Antitussive, antiasthmatic, hepatoprotective	41,42

Plant name	Family	Chemical constituents	Traditional uses	References
<i>Zingiber officinale</i>	Zingiberaceae	Gingerol, curcumin,	Catarrh, rheumatism, nervous diseases, gingivitis, toothache, asthma, stroke, constipation and diabetes. Anti-platelets, antioxidant, anti-tumor, anti-rhino viral, anti-hepatotoxicity, anti-arthritic and anti-inflammatory.	43,44,45,46,47
<i>Cucurbita pepo L. (pumpkin)</i>	Cucurbitaceae	Linoleic acid, oleic acid, Tocopherols, Sterols, proteins, peptides, polysaccharides, para amino benzoic acid and fixed oil. Pumpkin seed oil contains omega-6, omega-9, phytosterols, carotenoids, vitamin A, vitamin E, Palmitic acid	Anti-inflammatory, antiviral, analgesic, urinary disorders, anti-ulcer, antidiabetic and antioxidant.	48,4,50,51,52
<i>Citrus reticulata</i>	Rutaceae	Vitamin C, flavonoids, acids and volatile oils, hesperidin, rutin, naringin	Anticarcinogenic, anti-inflammatory, antioxidant, antimicrobial, hepatoprotective	53,54,55,56,57
<i>Petroselinum crispum oil</i>	Apiaceae	Apiin, luteolin, apiol, myristicin	Antibacterial, antioxidant, hepatoprotective	58,59,60
<i>Andrographis paniculata</i>	Acanthaceae	Flavonoid, phenol and glycosides	Antibacterial, antimalarial, antiviral, cardioprotective, antioxidant, anti-inflammatory, antidiabetic, antitumor activities.	61,62,63,64,65,66
<i>Silybum marianum</i>	Asteraceae	Silybin, silydianin, silychristin, betaine,	Anti-inflammatory	67,68
<i>Camellia sinensis</i>	Theaceae	Catechin, epicatechin, epigallocatechin, tannins, caffeine	Antioxidant, anticancer, lipid lowering activity, neuromuscular blocking action, Antiviral activity, antibacterial activity, antispasmodic activity, anticataract activity, antigenotoxic effect, anti-inflammatory, chemoprotective	108
<i>Acacia mellifera</i>	Fabaceae	Alkaloids, flavonoids, tannins, sterols, and saponins	Bowel problems, stomach trouble, cold, treatment for malaria and inflammation	2,21,46
<i>Adansonia digitata L.,</i>	Malvaceae	Terpenoids, flavonoids, steroids, vitamins, amino acids, carbohydrates and lipids	Immunostimulant, anti-inflammatory, analgesic, pesticide, antipyretic, febrifuge and astringent in the treatment of diarrhea	69,70,71
<i>Argemone Mexicana L.,</i>	Papaveraceae	Alkaloids as Berberine, protopine, sarguarine, optisine, and chelerytherine, the seed oil contains myristic, palmitic, oleic, linoleic acid, flavonoids, glycosides, sterols and phenolic compounds.	Analgesic, antispasmodic, hallucinogenic, sedative, tumors, warts, skin diseases, inflammations, rheumatism, jaundice, leprosy, piles, warm infestations and dysentery	76,77,78
<i>Anogeissus leiocarpus</i>	Combretaceae	Ellagic acid, Polyalcohol, sorbitol, terpenoids like α -amyrin, β -sitosterol, and traces of alkaloids	Parasitic diseases and dysentery	72,79,80,81
<i>Balanites aegyptiaca L.,</i>	Balanitaceae	Rotenone, yamogenin, steroidal sapogenin	Jaundice, liver disorders, spleen problems, the leaves and branches are used as a fumigant for rheumatism, fruits are used against constipation and antidiabetic	72,73,82,83,84
<i>Cannabis sativa L.</i>	Cannabaceae	Cannabinoids, Cannabidiol, Cannabinol	Spasmolytic, analgesic, hypnotic and to treat cholera, rabies, epilepsy, tetanus, rheumatism, anti-inflammatory	85
<i>Capparis deciduas</i>	Capparaceae	Alkaloids, saponins, flavonoids, tannins, sterols, cyanogenic, glycosides and coumarins	Jaundice, anti-rheumatic, diuretics, antigitout, anti-inflammatory, astringent, laxative, Stomachic, astringent and for skin diseases	86,72,73,87
<i>Combretum hartmannianum</i>	Combretaceae	Alkaloids, flavonoids, tannins, saponins, cyanogenic glycosides, sterols and coumarins	Barks and leaves extract are used for jaundice	69,72,88

Plant Name	Family	Chemical constituents	Traditional uses	References
<i>Khaya senegalensis</i>	Meliaceae	Saponins, tannins, alkaloids, glycosides, terpenoids, phenolic compounds and flavonoids, steroids	Diarrhea, dysentery and wound infections, fever and to treat syphilis, the bark used to treat jaundice, scorpion bites, allergies	79,89
<i>Kigelia Africana Lam</i>	Bignoniaceae	Verminosides and iridoids and series of polyphenols such as verbascoside	Bark is used to treat ulcer, pneumonia, malaria	69,72,90
<i>Lawsonia inermis L.,</i>	Lythraceae	Phenolic derivatives, xanthenes, naphthoquinone, coumarins, triterpenes, tannins, flavonoids, aliphatic components, sterols, gallic acid, mannitol, glucose, amino acids, trace elements and minerals.	Astringent, hypotensive, sedative and, headache, jaundice, leprosy and skin disease.	69,72,91
<i>Lepidium sativum L.,</i>	Brassicaceae	Triterpenes, alkaloids, tannins, flavonoids, coumarins and saponins	Gastrointestinal disorders, arthritis and inflammatory disorders, dysentery and diarrhea.	69,92
<i>Moringa oleifera Lam</i>	Moringiaceae	Alkaloids, saponins, flavonoids, tannins, sterols, glycosides and coumarins	Liver disease, lipid disorders, arthritis, inflammatory disorders, seeds used to clean water for drinking	69,72,93
<i>Nigella sativa L.,</i>	Ranunculaceae	Thymoquinone, thymohydroquinone, dithymoquinone, p-cymene, carvacrol, and 4-terpineol	Liver tonics, digestive, anti-inflammatory, immune-stimulant and remedy for jaundice anti-diabetics	94
<i>Occimum basilicum L.,</i>	Lamiaceae	Glycosides, gums, mucilage, proteins, amino acids, tannins, phenolic compounds, triterpenoids, sterols, saponins, flavones and flavonoids. Linalol methylchavicol, methylcinnamate, linolen, essential oil, rosmarinic acid, citral, eugenol and geraniol.	Jaundice, stomach complaints, fever, cough and gout diuretic, aphrodisiac, and anti-dysenteric. Seeds are used as demulcents.	73,95,96
<i>Phoenix dactylifera Linn</i>	Palmae	Carbohydrates, alkaloids, steroids, flavonoids, vitamins, tannins and phenolic acids	Sexual incapacity and weakness	72,97,98
<i>Raphanus sativus L.,</i>	Cruciferae	Flavonoids, saponins, coumarins, and alkaloids	Hepatoprotective, bacterial and viral infections, inflammation.	99
<i>Solanum nigrum L.</i>	Solanaceae	Gallic acid, catechin, Protocatechuic acid, caffeic acid, epicatechin, rutin and naringenin	Inflammatory disorders, rheumatism, swollen joints, hepatomegaly, splenomegaly, gonorrhoea, edema.	100,101
<i>Sterculia setigera Del.</i>	Sterculiaceae	Saponins, steroidal, sterols, flavonoids, Tannins, saponins, cardiac glycosides and anthraquinone	Jaundice, bilharzia	72,102,103
<i>Tamarindus indica L.</i>	Caesalpinaceae	Tannins, alkaloids, flavonoids, and saponins	To treat fever, postpartum remedy and measles	72,104,105,106

Conclusion

Liver is the largest metabolic organ in the human body. It has the function to detoxify toxic substances and synthesize useful biomolecules. So damage to liver may lead to grave consequences. This damage occurs due to chronic alcoholic abuse, viral hepatitis and inherited metabolic disease. Liver damage is associated with cellular necrosis, fibrosis, and increase in tissue lipid peroxidation and depletion in tissue glutathione level. Most of the hepatotoxic chemicals damage liver cells mainly by inducing lipid peroxidation and other oxidative damages. Polyphenols, flavonoids, terpenoids are natural antioxidants and the consumption of foods that contain these compounds in large quantities seems to play an important role in prophylaxis against many diseases.

REFERENCES

- A.Bilal, N. Jahan, A. Ahmed, Naaz Bilal, S. Habib, and S. Hajra, : Phytochemical and pharmacological studies on *Ocimum basilicum* linn- a review”. International Journal of Current Research and Review, vol.4, no.23, 2012.
- A.H.Arbab, M.K. Parvez, M.S. Al-Dosari t al., “Hepatoprotective and antiviral efficacy of *Acacia mellifera* leaves fractions against hepatitis B virus:. Biomed Research International, vol. 2015, Article 1D 929131, “, 10 page, 2015.
- A.I. Abuelgasim, E. A. Omer, and B. Elmahdi, “The effectiveness of *Nigella sativa* against liver damage in rats”, Research Journal of Medicinal Plants, vol.2, no.1, pp.43-47,2008.
- A.I. Abuelgasim, H. Nuha, and A. Mohammed, “Hepatoprotective effect of *Lepidium sativum* against carbontetrachloride induced damage in rats”, Research Journal of Animal & Veterinary Sciences, vol.3, pp.20-23,2008.
- A.Koch, P. Tamez, J. Pezzuto, and D. Soejarto, “Evaluation of plants used for antimalarial treatment by the Maasai of Kenya”. Journal of Ethnopharmacology, vol. 101,no, 1-3, pp. 95-99, 2005.
- Adi AH, Alturkmani HJ. Physiologically lucky: the role of medical physiology in modern medical education: an interview.Perspect Med Educ 2013; 2: 99–103.
- Aeschlbach R, Loliger J, Scott BC, Murcia A, Butler J, Halliwell B, et al. Antioxidant actions of thymol, carvacrol, 6-gingerol, Zingerone and hydroxytyrosol. Food chem. Toxicol 1994; 32:31-6.
- Ahsan R, Islam M, Bulbul JI, Musaddik A, Haque E. Hepatoprotective activity of methanol extract of some medicinal plants against carbon tetrachloride-induced hepatotoxicity in rats. Eur J Sci Res 2009; 37(2):302–10.
- Alemi M, Sabouni F, Sanjarian F, Haghbeen K, Ansari S. Anti- inflammatory effect of seeds and callus of *Nigella sativa* L. extracts on mix glial cells with regard to their thymoquinone content. AAPS PharmSciTech 2013; 14: 160-167.
- Al-Ghamdi MS. The anti-inflammatory, analgesic and antipyretic activity of *Nigella sativa*. J Ethnopharmacol 2001; 76: 45-48.
- Ali Esmail Al-Snafi, Chemical constituents and Pharmacological activities of Ammi majus and Ammi visnaga. A Review, International Journal of Pharmacy and Industrial Research, Vol-03(03) 2013, 257-265.
- Amagase H, Petesch BL, Matsuura H, Kasuga S, Itakura Y. Intake of garlic and its bioactive components. J Nutr 2001; 131: 955S-962S.
- Amin M, Pipelzadeh MH, Mehdinejad M, Rashidi I. An *in vivo* toxicological study upon shallomin, the active antimicrobial constitute of Persian shallot (*Allium hirtifolium*, Boiss) extract. Jundushapur J Nat Pharm Prod 2012; 7: 17-21.
- Asgari S, Setorki M, Rafieian-Kopaei M, Heidarian E, Shahinfard N, Ansari R, et al. Postprandial hypolipidemic and hypoglycemic effects of *Allium hirtifolium* and *Sesamum indicum* on hypercholesterolemic rabbits. Afr J Pharm Pharmacol 2012; 6: 1131-1135.

- Asif HM, Akram M, Usmanghani K, Akhtar N, Shah PA, Uzair M, et al. Monograph of *Apium graveolens* Linn. *J Med Plants Res* 2011; 5: 1494-1496.
- Awang DVC. Ginger. *Can Pharm J* 1992, 125: 309-11.
- Azadi HG, Riazi GH, Ghaffari SM, Ahmadian S, Khalife TJ. Effects of *Allium hirtifolium* (Iranian shallot) and its allicin on microtubule and cancer cell lines. *Afr J Biotechnol* 2009; 8: 5030-5037.
- B.A.Ahmed, "Hepatoprotective, antioxidant and antibacterial effects of *Ocimum basilicum*, *Anogeissus leiocarpus*, *Khaya senegalensis* and *Capparis decidua* against CCl₄ induced hepatotoxicity in rats". Sudan Academy of Sciences (SAS), Council of Bioscience, Advanced Technology and Environment, Sudan, 2015, PhD. thesis.
- B.A.Ahmed, A.H. Mohamed, S.A. Ali and A.I. Abuelgasim, "Hepatoprotective activity of ethanol extract of *Ocimum basilicum* against CCl₄ induced hepatotoxicity in albino rats", *Journal of Natural and Medical Sciences*, vol.16, no.2, pp. 11-18,2015.
- Bartalis J. Hepatoprotective activity of cucurbitacin [dissertation]. Brookings: South Dakota State University; 2005
- C.L.Priya ad K.V.B.Rao, "Ethnobotanical and current ethanopharmacological aspects *Argemone Mexicana* linn: an overview". *International Journal of Pharmaceutical Sciences and Research*, vol 3, no.7,p. 2143, 2012.
- Choudhary AK, Devi RS. Serum biochemical responses under oxidative stress of aspartame in Wistar albino rats. *Asian Pac J Trop Dis* 2014;4(1):S403–10.
- Chow JM, Shen SC, Huan SK, Lin HY, Chen YC. Quercetin, but not rutin and,quercetin, prevention of H₂O₂-induced apoptosis via anti-oxidant activity and heme oxygenase 1 gene expression in macrophages. *Biochem Pharmacol* 2005; 69(12): 1839-51.
- D.Chothani and H. Vaghasiya, " A review on *Balanites aegyptiaca* Del (desert date): phytochemical constituents, traditional uses and pharmacological activity". *Pharmacogony reviews*, vol, no.9, p. 55, 2011.
- D.Kabore, S.L.Hagretou, D.Brehima, C. Clarise, H.D. Mamoudou, and J.Mogens, "A review of Baobab (*Adansonia digitata*) products: effect of procesomg techniques, medicinal properties and uses." *African Journal of Food Science*, vol.5, no. 16, pp. 833-844,2011.
- Dua VK, Ojha VP, Roy R, Joshi BC, Valecha N, Devi CU, et al. Antimalarial activity of some xanthones isolated from the roots of *Andrographis paniculata*. *J Ethnopharmacol* 2004; 95:247–51.
- E.Musa, E.L. Badwi, E.A. Osman, and M.M.Dahab, "Hepatoprotective and toxicity assessment of *Cannabis sativa* seed oil in Albino rat". *International Journal of Clinical and Biological Sciences*, vol 1, pp.69-76, 2012.
- Elberry AA, Harraz FM, Ghareib SA, Gabr SA, Nagy AA, Abdel-Sattar E. Methanolic extract of *Marrubium vulgare* ameliorates hyperglycemia and dyslipidemia in streptozotocin-induced diabetic rats. *Int J Diabetes Mellit* 2011; 11: 1877-1878.
- Emim JA, Oliveira AB, Lapa AJ. Pharmacological evaluation of the anti-inflammatory activity of a citrus bioflavonoid, herperidin, and the isoflavonoids, duartin and claussequinone, in rats and mice. *J Pharm Pharmacol* 1994; 46(2): 118-22.

- F.W.Andrews, The Flowering plants of the Anglo-Egyptian Sudan, vol. 2, T.Buncle & Co Ltd, Arbroath, Scotland, 1952.
- F.W.Andrews, The Flowering Plants of the AngloEgyptian Sudan, vol.3, T.Buncle &Co. Ltd., Arbroath, Scotland, 1956.
- Fatehi M, Saleh TM, Fatehi-Hassanabad Z, Farrokhfal K, Jafarzadeh M, Davodi S. A pharmacological study on *Berberis vulgaris* fruit extract. *JEthnopharmacol* 2005; 102: 46-52.
- Fazal SS, Singla RK. Review on the pharmacognostical and pharmacological characterization of *Apium graveolens* Linn. *Indo Glob J Pharm Sci* 2012; 2: 36-42.
- G.B.E. El-Ghazali, S. M. El-Tohami, and E.E. Awatif “Medicinal plants of the Sudan” in Medicinal plants of Northern Kordofan, Omdurman Islamic University Printing and Publishing House, Omdurman, Sudan, 1997.
- G.E.B. El Ghazali, W.E. Abdalla, H.E. Khalid, M.M. Khalafalla and A.A. Hamad, Medicinal plants of Ingassana Area, National centre for research. Khartoum, Sudan, 2003.
- G.E.B. ELGhazali, Aromatic Plants of the Sudan, National centre for research. Khartoum University Press, Khartoum, Sudan, 2004.
- G.E.B.ELGhazali, M.S. Eltohami, and A.A.B. Elegami, Medicinal plants of the Sudan, Part III Medicinal plants of White Nile Province, National Centre for Research, Khartoum University Press, Khartoum, Sudan, 1994.
- Garg A, Garg S, Zaneveld LJ, Singla AK. Chemistry and Pharmacology of the citrus bioflavonoid hesperidin *Phyther Res* 2001; 15: 655-69.
- Gebhardt R. Antioxidative and protective properties of extracts from leaves of the artichoke (*Cynara scolymus* L.) against hydroperoxide- induced oxidative stress in cultured rat hepatocytes. *Toxicol Appl Pharmacol* 1997; 144: 279-286.
- Gião MS, Pereira CI, Fonseca SC, Pintado ME, Malcata FX. Effect of particle size upon the extent of extraction of antioxidant power from the plants *Agrimonia eupatoria*, *Salvia* sp. and *Satureja montana*. *Food Chem* 2009; 117: 412-416.
- Gossell-Williams M, Lyttle K, Clarke T, Gardner M, Simon O, Supplementation with pumpkin seed oil improves plasma lipid profile and cardiovascular outcomes of female non-ovariectomized and ovariectomized Sprague-Dawley rats. *Phyther Res* 2008; 22(7) 873-7.
- H.M. ELAmin, Trees & Shrubs of the Sudan, Itaca Press, Exeter, UK, 1990.
- H.M.E. Mohammed, “Pharmacological studies of selected plants with putative hepatoprotective activities”, Faculty of Pharmacy, Khartoum University, Khartoum, Sudan, 2008.
- Hadaruga DI, Hadaruga NG, Bandur GN, Ravis A, Costescu C, Ordodi VL, et al. *Berberis vulgaris* extract/ β cyclodextrin nanoparticles synthesis and characterization. *Rev Chim (Bucharest)* 2010; 61: 669-675.
- Hassan HA, Yousef MI. Ameliorating effect of chicory (*Cichorium intybus* L.)-supplemented diet against nitrosamine precursors-induced liver injury and oxidative stress in male rats. *Food Chem Toxicol* 2010; 48: 2163-2169.

- I.A.A. Abuowf and A.Abuowf, Hepatoprotective Activity of Date Palm (*Phoenix dactylifera*) Pollen Grains in Rats, University of Khartoum, Khartoum, Sudan, 2009.
- Imanshahidi M, Hosseinzadeh H. Pharmacological and therapeutic effects of *Berberis vulgaris* and its active constituent, berberine. *Phytother Res* 2008; 22: 999-1012.
- Jaeschke H, Gores GJ, Arthur IC, Jack AH, Pessayre D, Lemasters JJ. FORUM: mechanisms of hepatotoxicity. *Toxicol Sci* 2002;65:166–76.
- Kanter M, Coskun O, Budancamanak M. Hepatoprotective effects of *Nigella sativa* L and *Urtica dioica* L on lipid peroxidation, antioxidant enzyme systems and liver enzymes in carbon tetrachloride-treated rats. *World J Gastroenterol* 2005; 11: 6684-6688.
- Kshirsagar AD, Mohite R, Aggrawal AS, Suralkar UR. Hepatoprotective medicinal plants of ayurveda– a review. *Asian J Pharm Clin Res* 2011; 4: 1-8.
- Lantz RC, Chen GJ, Sarihan M, Solyom AM, Jolad SD, Timmermann BN. The effect of extracts from ginger rhizome on inflammatory mediator production. *Phytomedicine* 2007; 14; 123-8.
- Leelarasamee A, Trakulsomboon S, Maunwongyathi P, Somanabandhu A, Pidetcha P, Matrakool B, et al. Failure of *Phyllanthus amarus* to eradicate hepatitis B surface antigen from symptomless carriers. *Lancet* 1990; 335:1600–1.
- Liu J Lin H, McIntosh H. Genus *Phyllanthus* for chronic hepatitis B virus infection: a systematic review. *J viral Hepat* 2001; 8: 358-66.
- Lopez MG, Sanchez-Mendoza IR, Ochoa-Alejo N. Comparative study of volatile components and fatty acids of plants and in-vitro cultures of parsley (*Petroselinum crispum* (Mill) nym ex hill). *J Agric Food Chem* 1999; 47:3292–6.
- M.A. Mohamed, I.A. A. Mohamed, T.H. Tag El-Din, A. Hassan, and M.Hssan, : Hepatoprotective effect of *Adansonia digitata* L., (Baobab) fruits pulp extract on CCl₄ induced hepatotoxicity in rats:. *World Journal of Pharmaceutical Research*, vol 4, no.8, pp.368-377, 2015.
- M.Mohamed, I. Eldin, A. Mohammed, and Hassan, “Effects of *Lawsonia intermis* L., (Henna) leaves” methanolic extract on CCl₄ induced hepatotoxicity in rats:. *Journal of Intercultural Ethnopharmacology*, vol.5,no. 1, p.22,2016.
- M.Z.Zaruwa, N.I. Ibok et al., “Effects of *Stercula setigera* Del. Stem bark extract on haematological and biochemical parameters of wistar rats”, *Biochemistry Insight*, vol. 9, Article ID BCI. S36143, 2016.
- Makni M, Fetoui H, Gargouri NK, Garoui M, Jaber H, Maknni J, et al. Hypolipidemic and hepatoprotective effects of flax and pumpkin seed mixture rich in u-3 and u-6 fatty acids in hypercholesterolemic rats. *Food Chem Toxicol* 2008; 46: 3714-20.
- Mansi K, Abushoffa AM, Disi A, Aburjai T. Hypolipidemic effects of seed extract of celery (*Apium graveolens*) in rats. *Pharmacogn Mag* 2009; 5: 301-305.

- McBride A, Augustin KM, Nobbe J, Westervelt P. *Silybum marianum* (milk thistle) in the management and prevention of hepatotoxicity in a patient undergoing reinduction therapy for acute myelogenous leukemia. *J Oncol Pharm Pract* 2012; 18: 360-365.
- Mimica-Dukic N, Popovic M. Apiaceae species. A promising sources of pharmacologically active compounds and *Petroselinum crispum*, *Apium graveolens* and *Pastinaca sativa*. In: Govil JN, Singh VK, editors. Recent progress in medicinal plant species: phytopharmacology and therapeutic values III. Houston (TX): Studium Press LLC; 2007. p. 132–3.
- Molina-Montenegro MA, Atala C, Gianoli E. Phenotypic plasticity and performance of *Taraxacum officinale* (dandelion) in habitats of contrasting environmental heterogeneity. *Biol Invasions* 2010; 12: 2277-2284.
- Mumoli N, Cei M, Cosimi A. Drug-related hepatotoxicity. *N Engl J Med* 2006; 354(20):2191–3.
- Murkovic M, Hillibrand A, Winkler J, Pfannhauser W, Variability of vitamin E content in pumpkin seeds (*Curcubita pepo* L.) *Eur Food Res Technol* 1996; 202:275-8.
- N.H.S.H.Mohammed, A.I. Abelgasim, and A.H. Mohamed, “Protective effect of *Raphanus sativus* against carbon tetrachloride induced hepatotoxicity in Wistar albino rats”, *Journal of Pharmacology and Toxicology*, vol.3, no. 4, pp. 272-278,2008.
- Nagella P, Ahmad A, Kim SJ, Chung IM. Chemical composition, antioxidant activity and larvicidal effects of essential oil from leaves of *Apium graveolens*. *Immunopharmacol Immunotoxicol* 2012; 34: 205- 209.
- Okoh OO, Sadimenko AP, Asekun OT, Afolayan AJ. The effects of drying on the chemical components of essential oils of *Calendula officinalis* L. *Afr J Biotechnol* 2008; 7(10): 1500-1502.
- Preethi KC, Kuttan G, Kuttan R. Antioxidant potential of an extract of *Calendula officinalis* flowers *in vitro* and *in vivo*. *Pharm Biol* 2006; 44: 691-697.
- Punit R, Bhatt, Kajak B, Pandya, Navin R. Sheth, *Camellia Sinensis* (L): The Medicinal Beverage: A Review, *International Journal of Pharmaceutical Sciences Review and Research*, Volume 3, Issue 2, July-August 2010, Page no 6-9.
- R.A.M. Elhag and S.M.A. ElBadwi, “Hepatoprotective activity of *Solanum nigrum* extracts on chemically induced liver damage in rats”, *Journal of Veterinary Medicine and Animal Health*, vol.3, no.4. pp. 45-50, 2011.
- R.A.M. Elhag, “Hepatoprotective activity of two Sudanese plants on experimental liver damage”. Faculty of veterina.ry Science , Sudan, 2001.
- Rashid F, Ahmed R, Mahmood A, Ahmad Z, Bibi N, Kazmi SU. Flavonoid glycosides from *Prunus armeniaca* and the antibacterial activity of a crude extract. *Arch Pharm Res* 2007; 30: 932-937.
- Reichen J. The role of the sinusoidal endothelium in liver function. *News Physiol Sci* 1999;14:117.
- Ross & Wilson, Anne Waugh, Allison Grant, *Anatomy & Physiology in Health and Illness* 12th edition page no 308-310.

- S.A.Ali, “Hepatoprotective activity of selected sudanese plants on murine induced liver damage”, Faculty of Veterinary Science, Sudan, 2005.
- S.A.Ali, A.A. Gameel, A.H.Mohamed, and T.Hassan, “Hepatoprotective activity of Cpparis deciduas aqueous and methanolic stem extracts against carbon tetrachloride induced liver histological damage in rats”. *Journal of Pharmacology and Toxicology*, vol.6, no.1, pp. 62-68, 2011.
- S.A.Ali, T.H. Al-Amin, A.H. Mohamed, and A.A. Gameel, “Hepatoprotective activity of aqueous and methanolic extracts of Capparis decidua stems against carbon tetrachloride induced liver damage in rats”. *Journal of Pharmacology and Toxicology*, vol.4. no. 4, pp. 167-172, 2009.
- S.Adam, I. A. Idris, E. H. Abdelgadir, R.H. Ahmed, and E.E. Kamal. “Evaluation of hepatoprotective activity of Argemone Mexicana aerial part Extracts on CCl₄ induced liver damage in wistar rats.” *British Journal of Pharmacology and Toxicology*, vol.2, no.5, pp. 251-256,2011.
- S.Ali, A.H. Mohamed, G.E. Mohammed and A.A. Gameel, Toxicity of Balanites aegyptiaca seeds oil in wistar albino rats”, *SUST Journal of Agricultural and Veterinary Sciences*, vol.18, no1, 2017.
- S.Ali, A.H. Mohamed, A.A. Gameel, and T.Hassan, Protective Effect of Moringa Oleifera Leaves Aqueous Extract against Carbon tetrachloride induced Hepatotoxicity in rats, Science Congress Faculty of Veterinary Medicine, Assiut, Egypt, 2010.
- S.M.A. El Badwi, N.M. El Bagir, and A.E. Amin, “Protective effect of ethanolic extract of Tamarindus indica against CCl₄ induced liver damage in rats”, *Australian Journal of Basic and Applied Sciences*, vol.7. no.2, pp. 813-818, 2013.
- S.Sharma, I.Y.Adam, M. Marwa, and I.A.. Alhameed, “Kigelia Africana fruits” extracts and anti-hepatotoxic effects on male wistar rats liver destruction induced by CCl₄”. *Asian Journal of Medical Sciences*, vol.5, no.1, pp. 26-32, 2013.
- Sahpaz S, Garbacki N, Tits M, Bailleul F. Isolation and pharmacological activity of phenylpropanoid esters from Marrubium vulgare. *J Ethnopharmacol* 2002; 79: 389-392.
- Saller R, Meier R, Brignoli R. The use of silymarin in the treatment of liver diseases. *Drugs* 2001; 61:2035–63.
- Shaarawy SM, Tohamy AA, Elgendy SM, Abd Elmageed ZY, Bahnasy A, Mohamed MS, et al. Protective effects of garlic and silymarin on NDEA-induced rats hepatotoxicity. *Int J Biol Sci* 2009;5(6):549–57.
- Sheeja K, Shihab PK, Kuttan G. Antioxidant and anti- inflammatory activities of the plant *Andrographis paniculata* Nees. *Immunopharmacol Immunotoxicol* 2006; 28:129–40.
- Shukla B, Visen P, Patnaik GK, Dhawan BN. Choloretic effect of picroliv, the hepatoprotective principle of *Picrorhiza kurroa*. *Planta Med* 1991; 57: 29-33.
- Singh MK, Sahu P, Nagori K, Dewangan D, Kumar T, Alexander A, et al. Organoleptic properties *in-vitro* and *in-vivo* pharmacological activities of *Calendula officinalis* Linn: an over review. *J Chem Pharm Res* 2011; 3: 655-663.

- Stevenson DG, Eller FJ, Wang L, Jane JL Wang T, Inglett GE. Oil and Tocopherol content and composition of pumpkin seed oil in 12 cultivars J Agric Food Chem 2007; 55 (10): 4005-13.
- Subramaniam S, Khan HBH, Elumalai N, Lakshmi SYS. Hepatoprotective effect of ethanolic extract of whole plant of *Andrographis paniculata* against CCl₄-induced hepatotoxicity in rats. Comp Clin Path 2015;24:1-7.
- Subramaniam S, Khan HBH, Elumalai N, Lakshmi SYS. Hepatoprotective effect of ethanolic extract of whole plant of *Andrographis paniculata* against CCl₄-induced hepatotoxicity in rats. Comp Clin Path 2015; 24:1-7.
- T.H. Abdelrahman, “ Hepatoprotective activity of ethanolic and ethyl acetate extracts of *Sterculia Setigera* against Carbon tetrachloride induced hepatotoxicity in albino rats”, Mediterranean Journal of Biosciences, vol.1. no.3, pp. 114-119, 2016.
- T.Okpekon, S. Yolou, C. Gleye et al., “Antiparasitic activities of medicinal plants used in Ivory coast”. Journal of Ethnopharmacology, vol 90, no.1, pp. 91-97,2004.
- T.S.Sourabie, H.M. Kone, J. B. Nikiema, O.G. Nacoulma, and I.P. Guissou “Evaluation of the antihepatotoxic effect of *Argemone Mexicana* leaf extracts against CCl₄-induced hepatic injury in rats.” International Journal of Biological and Chemical Sciences, vol.3, no.6, 2009.
- Tabassum N, Shah MY, Qazi MA, Shah A. Prophylactic activity of extract of *Taraxacum officinale* Weber. against hepatocellular injury induced in mice. *Pharmacologyonline* 2010; 2: 344-352.
- Tanaka T, Kohno H, Tsukio Y, Honjo S, Tanino M, Miyake M. citrus limonoids obacunone and limonin inhibit azoxymethane-induced colon carcinogenesis in rats. *Biofactors* 2000; 13: 213-18.
- Tapsell LC, Hemphill I, Cobiac L, Patch CS, Sullivan DR, Fenech M, et al. Health benefits of herbs and spices: the past, the present, the future. *MedJ Aust* 2006; 185(4); S4-24.
- Tatiya AU, Surana SJ, Sutar MP, Gamit NH. Hepatoprotective effect of poly herbal formulation against various hepatotoxic agents in rats. *Pharmacognosy Res* 2012; 4: 50-56.
- Touloupakis E, Ghanotakis DF. Nutraceutical use of garlic sulfur- containing compounds. *Adv Exp Med Biol* 2010; 698: 110-121.
- Vamanu E, Vamanu A, Nita S, Colceriu S. Antioxidant and antimicrobial activities of ethanol extracts of *Cynara scolymus* (*Cynarae folium*, Asteraceae family). *Trop J Pharm Res* 2011; 10: 777-783.
- Wang WH, Wang AM. Studies of commonly used traditional medicine-ginger. *Zhongguo Zhong Yao Za Zhi* 2005; 30 1569-73.
- Wang YC, Chuang YC, Hsu HW. The flavonoid, carotenoid and pectin content in peels of citrus cultivated in Taiwan. *Food Chem* 2008; 106:277-84.
- Wiat C, Kumar K, Yusof MY, Hamimah H, Fauzi ZM, Sulaiman M. Antiviral properties of ent-labdene diterpenes of *Andrographis paniculata* Nees, inhibitors of herpes simplex virus type 1. *Phytother Res* 2005; 19:1069-70.

- Wong PYY, Kitts DD. Studies on the dual antioxidant and anti- bacterial properties of parsley (*Petroselinum crispum*) and cilantro (*Coriandrum sativum*) extracts. *Food Chem* 2006; 97:505–15.
- Wu L, Chen Y. Protective effect of celery root against acute liver injury by CCl₄. *West China J Pharm Sci* 2008; doi: 10.3969/j.issn.1006-0103.2008.04.013.
- Y.J.Hong, F.A.Tomas-Barberan, A.A. Kader, and A.E.Mitchell, “ The flavonoid glycosides and procyanidin composition of deglet noor dates (*Phoenix dactylifera*)”, *Journal of Agricultural and Food Chemistry*, vol.54, no.6,pp. 2405-2411.2006.
- Yildiz F, Coban S, Terzi A, Ates M, Aksoy N, Cakir H, et al. *Nigella sativa* relieves the deleterious effects of ischemia reperfusion injury on liver. *World J Gastroenterol* 2008; 14: 5204-5209.
- Zhao F, He EQ, Wang L, Liu K. Anti-tumor activities of andrographolide, a diterpene from *Andrographis paniculata*, by inducing apoptosis and inhibiting VEGF level. *J Asian Nat Prod Res* 2008; 10:467–73.