


International Journal of  
**Biology**  
(IJB)

**INFLUENCE OF INTRAPERITONEAL ADMINISTRATION  
OF CERIUM OXIDE ON BLOOD PARAMETERS AND LIPID  
PROFILE IN MICE (*MUS MUSCULUS*)**

Samaher Sabri Hammid and Sami Jeber Al-Maliky



## INFLUENCE OF INTRAPERITONEAL ADMINISTRATION OF CERIUM OXIDE ON BLOOD PARAMETERS AND LIPID PROFILE IN MICE (*MUS MUSCULUS*)

<sup>1</sup>Samaher Sabri Hammid

<sup>1</sup>Department of medical sciences college of nursing –university of Basrah-Iraq

<sup>2</sup>Sami Jeber Al-Maliky

<sup>2</sup>Prof. College of Education for pure sciences - university Basrah University-Iraq

### Abstract

**Purpose:** White laboratory mice *Mus musculus* L. strain BALB / C were used to study the effects cerium oxide on the hematological parameters and lipid profile values.

**Results:** Results showed a significant decrease ( $p \leq 0.05$ ) in the RBCs' hemoglobin Hb 'and PCV' parameters as well as the presence of a significant in the mean erythrocyte MCV volume in the two doses (2.5'and .3.75) micro liters / kg ', while there was no significant difference in hemoglobin (MCH) and the mean concentration of hemoglobin in red blood cells (MCHC) when compared with the control group, while the two doses did not differ significantly in PCV, Hb, RBCs, while the two doses differed significantly between them in MCV. There is significant difference in the level of triglycerides (TG, LDL) and HDL (compared with the control group) for both doses, and it was found that there was a significant difference between the low and high dose (2.'3.75) micro liters / kg in the HDL level. TG, LDL) compared with the control group at the probability level ( $P \leq 0.05$ ), while there was no difference between the high and low dose in the level of cholesterol and (VLDL).

**Key words:** *Cerium oxide, liver enzymes, Kidney, rats*

## INTRODUCTION

The widespread interest in nanoparticles comes from the small particle size resulting in novel properties such as 'enhanced chemical interaction', optical behavior and large surface (1,2,3). It is already used in many applications including household products, equipment, food, medicine, optics, sun lotions, cosmetics, paints, textiles, electronics, and bioremediation production (4, 5).

The increasingly widespread use and interest shown by the industry about NPs and consequently on nanomaterials 'increased distribution resulting in increased public and occupational exposure' and thus raised questions about the potential impact of these new materials on the environment and human health (6). Cerium oxide is an example of nanoparticles which are the basis as additive sources of fuel to enhance combustion efficiency and clean the engine 'and reduce fuel consumption and emissions to the environment (7).

Cerium oxide nanoparticles (CNPs) that consist of cerium atoms bonded to oxygen atoms' have been used in commercial, industrial and biological applications (8). Consequently the present study carried out to investigate the effect of cerium oxide on blood and lipids profile parameters

## MATERIAL AND METHODS

The current study was conducted on white laboratory mice *Mus musculus* L. strain BALB / C that was obtained from (Center for Drug Control - College of Science, Dhi Qar University). The mice were raised in the animal house of the Department of biology department - College of Education in Pure Sciences - University of Basra under Controlled conditions with a temperature of 20-25 ° C and a lighting cycle of 12 hours of light and 12 hours of darkness (AL-Malki, 2000). The mice were placed in plastic cages of standard sizes 30 X 12 X 11 cm manufactured by North Kint Plastic Kint.UK. . Sawdust was used as a bed and it was replaced weekly.

Male mice were divided, 10-12 weeks old, with a weight of 20-25 grams for each test mice divided into three groups consisting of eight mice each as follows (1) The control group Laboratory mice were injected with 1 ml of Normal saline, (2) Treated group, were injected with 1 ml of cerium oxide at a concentration of 2.5 l of the raw material to 997.5 l of the physiological solution. Equivalent to 0.05 mg / 100 body weight. The second group were injected with 1 ml of cerium oxide at a concentration of 3.75 of the raw material to 996.25 l of physiological solution, equivalent to 0.05 ml / 100 grams of body weight. Duration of the Injection period were 15 and 30 days, with one injection every two days, for 15, 30 days. Effective Material is a substance of cerium oxide found in the form of a concentrated liquid solution in a package volume of 100 milliliters containing the active substance cerium oxide at a concentration of 20%, produced by SIGMA (USA). Draw blood from the heart directly by the cardiac puncher using a 1 ml disposable syringe, put the blood in tubes containing EDTA for blood parameters tests. As for the blood for the purposes of biochemical tests, the blood was placed in glass tubes free of anticoagulant. It contains a gelatinous substance tube and left for 10-15 minutes to coagulate, then the tubes were placed in a Centrifuge for 10 minutes at 3500 rpm in order to obtain an adequate amount of blood serum, then the serum was drawn from the coagulated blood and kept in special tubes with a at -20 ° C until the biochemical tests are performed. Physiological parameters were measured using a hematology analyzer and spectrophotometer for lipids measurement. An analysis of variance one way using Statistical Package for Social Sciences (SPSS, V.11) was used.

## RESULTS AND DISCUSSION

Table (1) showed the results of the effect of cerium oxide on the hematological parameters of male laboratory mice. Results showed a significant decrease at the probability level ( $p \leq 0.05$ ) in the RBCs' hemoglobin Hb 'and PCV' as well as the presence of a significant decrease ( $P \leq 0.05$ ) in the mean erythrocyte MCV volume in the two doses (2.5, 3.75) micro liters / kg 'while there was no significant difference in the rate of hemoglobin (MCH) and the mean concentration of hemoglobin in red blood cells (MCHC) when compared with the control group, while the two doses did not differ significantly between them in the rate of PCV, Hb, RBCs, while the two doses differed significantly between them in the rate of MCV. our results showed agreement with (9) who revealed that diets supplemented with CeO, particularly at 200 ppm concentration, is capable of improving body weight gain, daily dry matter intake and feed conversion ratio in rabbits without any detrimental effect on the blood parameters indicating good health status of the animals.

**Table ( 1 ) The effect of cerium oxide on the mean of some blood parameters in male laboratory mice For fifteen days (n = 7) (Mean  $\pm$  SE).**

\*I different from control  $p < 0.05$

A study by (10) demonstrated the effect of toxicity of cerium oxide nanoparticles on blood cells in male mice, where the mean total hematocrit and average red blood cell (RBC) count were observed in all experimental groups; but white blood cells (WBC) compared to the group Controls' only have linear increment

Parameters Group	RBC count Cell/ml $\times 10^6$	Hb g/dL	PCV %	MCV ft	MCH pg	MCHC %
Control 1ml saline	56.7 $\pm$ 0.081	13.457 $\pm$ 0.287	45.00 $\pm$ 0.899	61.285 $\pm$ 1.215	17.971 $\pm$ 0.229	30.328 $\pm$ 0.379
T1 1ml (2.5 uL/100g CeO	L 4.248 $\pm$ 0.404	L 6.921 $\pm$ 0.393	L 23.771 $\pm$ 1.678	l 50.5000 $\pm$ 2.592	16.571 $\pm$ 0.666	30.615 $\pm$ 0.445
T2 ml 3.7 (uL/100g CeO	L 4.287 $\pm$ 0.609	L 7.957 $\pm$ 1.155	L 22.773 $\pm$ 2.890	L 55.959 $\pm$ 2.283	18.128 $\pm$ 0.671	31.614 $\pm$ 1.082



**Table ( 2 ) The effect of cerium oxide on the mean of some blood parameters in male laboratory mice For thirty days (n = 7) (Mean  $\pm$  SE).**

Parameters Group	RBC count Cell/mlX10 <sup>6</sup>	Hb g/dL	PCV %	MCV ft	MCH pg	MCHC %
<b>Control 1ml saline</b>	56814 $\pm$ 06485	13.385 $\pm$ 0.318	45.142 $\pm$ 0.769	59.214 $\pm$ 0.634	17.728 $\pm$ 0.357	30.214 $\pm$ 0.219
<b>T1 1ml (2.5 <math>\mu</math>L/100g) CeO</b>	1 4.400 $\pm$ 0.691	1 5.714 $\pm$ 0.449	1 20.071 $\pm$ 0.889	1 46.628 $\pm$ 1.107	1 16.185 $\pm$ 0.670	1 856.74 $\pm$ 0.6318
<b>T2 3.7 ml (3.7 <math>\mu</math>L/100g) CeO</b>	1 4.171 $\pm$ 0.494	1 5.285 $\pm$ 0.401	1 24.285 $\pm$ 3.746	1 44.785 $\pm$ 1.757	1 16.814 $\pm$ 1.165	1 30.616 $\pm$ 1.084

\*I different from control  $p < 0.05$ 

Low Hb and PCV indicate presence of anemia, (11) insure the founding of the recent study, he found that treated group showed normocytic normochromic anemia at the 7th day and microcytic hypochromic anemia at the 15th day.

The results in Table ( 2 ) showed a significant decrease in the red blood cells (RBCs), the rate of hemoglobinemia (Hb), the volume of compressed blood PCV, and the average red blood cell volume (MCV) in the two doses (2.5-3.75) micro liters / kg show a decrease 'at a probability level (0.05)  $P \leq$  when compared with the control group, while the two doses did not show a significant difference in the rate of hemoglobin MCH and the mean hemoglobin concentration in the red blood cell MCHC when compared with the control group, but at the level of the two doses, exhibited significantly differ in RBCs, Hb, PCV, MCV, MCH and MCHC.

The findings of (12) exhibit that, although CNPs did not significantly perturb the native conformation of Hb, they can stimulate some cellular adverse effects at high concentrations

The current study results showed a significant increase in the level of total cholesterol and HDL and LDL' for male laboratory mice injected with a high dose (3.75) micro liters / kg of cerium oxide at a probability level ( $p \leq 0.05$ ) compared with the control group 'while there was no significant difference. In the level of triglycerides (TG) and VLDL) compared with the control group at the probability level ( $P \leq 0.05$ ) for both doses, It was also found that there was a significant difference between the low and high dose (2.5-3.75) micro liters / kg in the LDL level compared with the control group at the level of ( $P \leq 0.05$ ) 'while there was no difference between high and low dose in cholesterol and (LDL)'.

The results also showed that there was a significant difference between the low dose and the high dose compared with the control group 'in the level of HDL and the presence of a significant difference between the two doses when compared to them' as shown in the table ( 3)

**Table (3) The effect of cerium oxide on lipid profile of male laboratory mice for fifteen days(n = 7) (Mean  $\pm$  standard error**\*I different from control  $p < 0.05$

The results of the current study showed a significant increase in the level of total cholesterol and HDL 'LDL' for male laboratory mice injected with a high dose (3.75) micro liters / kg 'of cerium oxide at a probability level ( $p \leq 0.05$ ) compared with the control group' while it was found that there is Significant difference in the level of triglycerides (TG, LDL) and HDL (compared with the control group) at the probability level ( $P \leq 0.05$ ) for both doses, and it was found that there was a significant difference between the low and high dose (2.5'3.75) micro liters / kg in the HDL level. TG, LDL) compared with the control group at the probability level ( $P \leq 0.05$ ), while there was no difference between the high and low dose in the level of cholesterol and (VLDL), as shown in the table.

**Table (4) The effect of cerium oxide on lipid profile of male laboratory mice for thirty days (n = 7) (rate  $\pm$  standard error)**

\*I different from control  $p < 0.05$

Plasma HDL-C concentrations are not an appropriate marker of vascular effects of HDL and therefore do not represent a reliable therapeutic target (13). Recent studies have focused on measuring other indices of HDL, such as the functionality, size, or number of HDLparticle. Of all the indices, two markers associated with HDL functionality, i.e., HDL-LOOH and HDL-TG, are clinically available markers.

A lipid panel including these two markers is expected to serve as an initial screening

Parameters Group	Cholesterol mg/dL	TG g/dL mg/dL	HDL mg/Dl	LDL mg/dL	VLDL mg/dL
Control 1ml saline	117.476 $\pm$ 1.674	131.909 $\pm$ 12.922	77.585 $\pm$ 958	23.454 $\pm$ 742	26.381 $\pm$ 2.584
T11 ml (2.5 uL/100g ) CeO	1 133.397 $\pm$ 1 3.751	1 111.360 $\pm$ 1 56.58	1 87.260 $\pm$ 1 2.317	1 31.027 $\pm$ 1 817	1 22.271 $\pm$ 1 1.938
T2 ml 3.7 (uL/100g ) CeO	1 137.783 $\pm$ 1 2.136	1 115.325 $\pm$ 1 9.340	1 93.541 $\pm$ 1 1.689	1 33.505 $\pm$ 1 1.093	1 23.065 $\pm$ 1 1.868

procedure for predicting approximate risks of cardiovascular disease. (14).

That administration of sodium selenite and a combination of sodium selenite and CeO<sub>2</sub> nanoparticles significantly reduced ( $P < 0.05$ ) plasma LDL compared to the diabetic group(15). The nano and metal forms of of CeO<sub>2</sub> did not cause significant change. Treatment with CeO<sub>2</sub>NPs was able to reduce the size and content of hepatocyte lipid droplets, the hepatic concentration of triglyceride(16).

## Conclusion

The present study concluded that different doses and periods of intraperitoneal administration of cerium oxide nanoparticle in mice effect negatively blood parameters while elevated lipid profile,

## REFERENCES

- (1) Ju-Nam, and J. R. Lead. Manufactured nanoparticles: An overview of their chemistry, interactions and potential environmental implications. *The Science of the Total Environment* .2008-400 (1-3) 396-414
- (2) Gaiser B. ; Teresa F Fernandes, Mark A Jepson, Jamie R Lead, Charles R Tyler, Mohammed Baalousha, Anamika Biswas, Graham J Britton, Paula A Cole, Blair D Johnston, Yon Ju-Nam, Philipp Rosenkranz, Tessa M Scown, Vicki Stone . Interspecies comparisons on the uptake and toxicity of silver and cerium dioxide nanoparticles. *Environ Toxicol Chem.*2012) ;31(1):144-54
- (3) Xia D, *et al.*,. Structural analysis of cytochrome bc1 complexes: implications to the mechanism of function. *Biochim Biophys Acta* .2013)1827(11-12):1278-94.
- (4)Shaw P., Sharp W, Jonathan B. Attention-deficit/hyperactivity disorder is characterized by a delay in cortical maturation. *National Academy of Sciences.*2007- 104(49):19649-54
- (5) Jessica T. Dahle and Yuji Arai .Environmental Geochemistry of Cerium: Applications and Toxicology of Cerium Oxide Nanoparticles. *IJERPH 2015-Volume 12 Issue 2* 10.3390
- (6) Dreher A.(2004)The Influence of IMF Programs on the Re-election of Debtor Governments.*Economic and politics Volume16, Issue1Pages* 53-76.
- (7) Park, J., Park, S., Seryun Kim, S., Jaeyoung C., Ahn, J., Seungmin L., Sunggon K. An informatics pipeline supporting phylogenomic analysis of fungal transcription factors .*Bioinformatics*,2008- Volume 24, Issue 7, , Pages 1024–1025,
- (8)Reeda, C Daniel , Christopher K. Algarb , Julie A. Huberb , and Gregory J. Dicka Genecentric approach to integrating .*environmental genomics and biogeochemical models* .February 4, 2014 . vol. 111 . no. 5 : 1879–1884
- (9)Adua OA and Akinmuyisitana IW (2013) Growth performance and blood profile of female rabbits fed dietary Cerium Oxide .*Journal of Bio-Science*. Vol 21 .
- (10)Massod Negahdary, A Asadi, S Mehrtashfar, M Imandar, H Akbari-Dastjerdi A biosensor for determination of H<sub>2</sub>O<sub>2</sub> by use of HRP enzyme and modified CPE with ZnO NPsInternational .*Journal of Electrochemical Science.*2012- 7 (6), 5185-5194
- (11)Ali Alkaladi a,d , Nasr A.M. Nasr El-Deen b , Mohamed Afifi a,c,d , Osama A. Abu Zinadah a, (2015) Hematological and biochemical investigations on the effect of vitamin E and C on *Oreochromis niloticus* exposed to zinc oxide nanoparticles *Saudi Journal of Biological Sciences*Volume 22, Issue 5, Pages 556-563
- (12) Neda Eskandari, , Sanaz Nikpur,2 Ghazal Ghasrahmad, Farnoosh Attar,*et al.*,(2018)Biophysical, docking, and cellular studies on the effects of cerium oxide nanoparticles on blood components: in vitro. *Int J Nanomedicine*. 2018; 13: 4575–4589
- (13)Winfried März, Marcus E. Kleber, Hubert Scharnagl, Timotheus Speer, Stephen Zewinger, Andreas Ritsch, Klaus G. Parhofer, Arnold von Eckardstein, Ulf Landmesser, and Ulrich Laufs (2017)HDL cholesterol:reappraisal of its clinical relevance. *Clin Res Cardiol*. 2017; 106(9): 663–675.

- (14) Fumiaki Ito and Tomoyuki Ito .High-Density Lipoprotein (HDL) Triglyceride and Oxidized HDL: New Lipid Biomarkers of Lipoprotein-Related Atherosclerotic Cardiovascular Disease. *Antioxidants (Basel)*.;2020- 9(5): 362.
- (15) Nazila Pourkhalili, Asieh Hosseini, Amir Nili-Ahmadabadi, Shokoufeh Hassani, Mohsen Pakzad, Maryam Baeri, Azadeh Mohammadirad, and Mohammad Abdollahi(2011)(Biochemical and cellular evidence of the benefit of a combination of cerium oxide nanoparticles and selenium to diabetic rats. *World J Diabetes* 15; 2(11): 204–210.
- (16) Silvia Carvajal , Meritxell Perramón , Denise Oró , Eudald Casals , Guillermo FernándezVaro , Gregori Casals , Marina Parra , Bernardino González de la Presa , Jordi Ribera , Óscar Pastor , Manuel Morales-Ruíz , Víctor Puentes , Wladimiro Jiménez (2019)Cerium oxide nanoparticles display antilipogenic effect in rats with nonalcoholic fatty liver disease. *Sci Rep.* 6;9(1):12848